

**POISON**

KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# NEOVEDOR

## Pour-On Endectocide

ACTIVE CONSTITUENT: 5 mg/mL DORAMECTIN

For the treatment and control of doramectin sensitive internal and external parasites of cattle.

For the treatment and persistent control of gastrointestinal roundworms, lungworms and buffalo fly in cattle. Control of eye worms, sucking and biting lice, mange mites and cattle ticks. Highly effective in the treatment and control of the following species:

**Gastrointestinal roundworms (adults and forth stage larvae)**

*Ostertagia ostertagi* - Small brown stomach worm  
*Haemonchus placei* - Barber's pole worm  
*Trichostrongylus axei* - Stomach hair worm  
*T. colubriformis* - Black scour worm  
*Cooperia spp.* - Small intestinal worm  
*Cooperia oncophora*  
*Bunostomum phlebotomum* - Hook worm  
*Oesophagostomum radiatum* - Nodule worm  
*Nematodirus spathiger* - Thin necked intestinal worm  
*Thelazia spp.* - Eye worm  
*Thichuris spp.* - Whipworm

**Adult and Immature Lungworm**

*Dictyocaulus viviparus*

**Sucking Lice**

*Linognathus vituli*  
*Haematopinus eurysternus*  
*Solenopotes capillatus*

**Biting Lice**

*Bovicola bovis*

**Buffalo Fly**

*Haematobia irritans exigua*

**Cattle Tick**

*Boophilus microplus*

**Mange Mites**

*Chorioptes bovis*  
*Sarcoptes scabiei var. bovis*

When applied as directed this product can persistently protect against various gastrointestinal roundworms for 21 to 35 days; against lungworms for up to 28 days and against buffalo flies for up to 14 days.

Extensive studies have also demonstrated the following levels of persistent protection against reinfection from important worm species:

*Ostertagia ostertagi* - Up to 35 days  
*Haemonchus placei* - Up to 35 days  
*Trichostrongylus axei* - Up to 35 days  
*Cooperia spp.* - Up to 35 days  
*Cooperia oncophora* - Up to 21 days  
*Dictyocaulus viviparus* - Up to 28 days  
*Oesophagostomum radiatum* - Up to 21 days  
*Bunostomum phlebotomum* - Up to 21 days

**DIRECTIONS FOR USE:****Restraints:**

Animals must NOT be repeat treated within 56 days (8 weeks) of a previous treatment.

**Precautions:**

For best results, do not treat cattle when their hide or hair is wet.

Do not apply to areas of skin which are contaminated with mud or manure.

Worm parasites may develop resistance to any anthelmintic. If this is suspected contact your local veterinary practitioner or advise the manufacturer.

**Dosage and Administration:****For external use only.**

Apply topically along the mid-line of the back in a narrow strip between the withers and tailhead.

Use the dose rate, according to the weight of the animal in the weight/dose table. The dose rate should be based on the heaviest member in the weight range. Cattle should be weighed with either scales or a weighband.

**DOSE RATE:** 1mL per 10 kg bodyweight.

| WEIGHT (kg) | DOSE (mL) |
|-------------|-----------|
| 40 - 50     | 5         |
| 51 - 75     | 7.5       |
| 76 - 100    | 10        |
| 101 - 150   | 15        |
| 151 - 200   | 20        |
| 201 - 250   | 25        |
| 251 - 300   | 30        |
| 301 - 350   | 35        |
| 351 - 400   | 40        |
| 401 - 450   | 45        |
| 451 - 500   | 50        |
| 501 - 550   | 55        |
| 551 - 600   | 60        |
| 601 - 650   | 65        |

\*Cattle whose weight exceeds 650 kg should be dosed at 1 mL per 10 kg bodyweight.

Treatment of very young animals less than 40 kg bodyweight is generally not warranted.

To obtain optimal benefit from the persistent-action, it is recommended that cattle which are set-stocked should be treated twice at eight week intervals. By using in this way, Neovedor Pour-On Endectocide will reduce the build-up of infective larvae on pasture and protect such as animals from parasitic gastro-enteritis and parasitic bronchitis by reducing pasture larval challenge.

To obtain optimal benefit from the persistent-action against internal parasites, it is recommended that cattle which are set-stocked should be treated twice at eight week intervals.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL DIRECTIONS:**

**WITHHOLDING PERIODS:**

**MEAT - DO NOT USE** less than 42 days before slaughter for human consumption.

**MILK - NIL**

**TRADE ADVICE:**

**Export Slaughter Interval:** Do not slaughter cattle for export within 42 days of treatment.

**SAFETY DIRECTIONS:**

Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with skin and eyes. When using the product wear face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after

use. After each day's use, wash face shield or goggles.

Use only in well ventilated surroundings.

**Flammable -** Keep away from heat, sparks, open flame and other sources of ignition.

**FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

**ENVIRONMENTAL PROTECTION:**

Doramectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

When used according to label instructions, the product is not likely to have an unintended effect that is harmful to the environment or have an overall adverse impact on dung beetle populations. Increased mortality and impaired development of larvae may be expected, but for a limited period after treatment.

**DISPOSAL:**

Triple-rinse container. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

**STORAGE:**

Store at temperatures below 30°C (Room Temperature). Protect from direct sunlight.

APVMA Approval No.: 83969 / 117495

**NEOVE PHARMA AUSTRALIA PTY LTD**

303c, Level 3, 276 Pitt Street, Sydney NSW 2000  
ACN 140 367 442

Customer Information Line: 1300 052 066

[www.neovepharma.com.au](http://www.neovepharma.com.au)

[www.vetproductsforfarmers.com.au](http://www.vetproductsforfarmers.com.au)

**Manufactured in Australia**